Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Marsupella sphacelata (Gieseke) Dum. Mt. Mansfield, Evans, D. C. Eaton.

Mylia anomola (Hook.) S. F. Gray. Jerico, Evans; Willoughby, Miss Lorenz.

**Nardia crenulata* (Smith) Lindb. Newfane, M. A. Howe, A. J. G.

N. hyalina (Lyell) Carrington. Newfane, M. A. Howe.

Scapania irrigua (Nees) Dum. Jerico, Evans.

S. paludosa (C. Muell.) Fib. Mt. Mansfield, Evans.

Sphenolobus exsectaeformis (Breidl.) Steph. Jerico, Evans.

* *S. exsectus* (Schmid.) Steph. Willoughby, Miss Lorenz.

**Anthoceros leavis* L., Newfane, Burlington, A. J. G.

**A. punctatus* L., Newfane, A. J. G.

Notothylas orbicularis (Schwein.) Sulliv. Jerico, Evans

TWO CHANGES OF NAME.

JOHN M. HOLZINGER.

RHACOMITRIUM FLETTII. Mr. Cardot after carefully examining a part of the type of this species agrees that it is new, but he considers it should be a Grimmia, being closely related to *Grimmia torquata* Grev., *G. prolifera* C. M. & K., and *G. tortifolia* Kindb., a group of species belonging to the subgenus Rhabdogrimmia of Limpricht, I accept my friends suggestion after looking over the matter, and *Rhacomitrium Flettii* may hereafter be known as **Grimmia Flettii** (Holz.) Card. See BRYOLOGIST **7**:3, 1904.

BRYUM SQUARROSUM Kindb. in Hedwigia, p. 66,1896. This name is not tenable being a synonym for *Paludella squarrcsa.* I propose for this Kindbergian plant **Bryum Baileyi** (Synonym *Bryum squarrosum* Kindb., non Linn.) in recognition of my very efficient bryological friend, Dr. John W. Bailey, of Seattle, Wash., who collected this plant at Lake Union near Seattle, in quantity for my Acrocarpi. I am indebted to Mr. Jules Cardot for its final determination. The nearest relative of this West American species is *Bryum Donianum* of the section Eubryum. But this has a thickened margin and smaller leaf cells. *Bryum Baileyi* has the margin of *one* cell layer, its leaf cells in the lower part of the leaf measure 24 by 120μ. The spores are smooth and measure 10 by 14μ, while Limpricht records the size of spores of *B. Donianum* as 9–10μ.

CURRENT LITERATURE.

Musci Norvegiæ Borealis by Dr. I. Hagen.

JOHN M. HOLZINGER.

The third and concluding part of this work on the mosses of Northern Norway, covering pages 241 to 382 and accompanied by twenty-four pages of Introduction appeared in February, 1905. The two plates, figure distinctive features of *Gyroweisia tenuis compacta*, *Distichium Hagenii*, *Encalypta mutica*, *Timmia Norvegica*, *T. elegans*, *Polytrichum inconstans* and *Hypnum fastigiatum mitodes*.